

AD-A271 849

UNCLASSIFIED
SECURITY CLASSIFICATION OF THIS PAGE

P. 104 & 105 (2)

Form Approved
OMB No. 0704-0188

REPORT

1a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION / AVAILABILITY OF REPORT Approved for public release; distribution is unlimited.		
2b. DECLASSIFICATION / DOWNGRADING SCHEDULE			4. MONITORING ORGANIZATION REPORT NUMBER(S)		
4. PERFORMING ORGANIZATION REPORT NUMBER(S)			5. MONITORING ORGANIZATION REPORT NUMBER(S)		
6a. NAME OF PERFORMING ORGANIZATION Smithsonian Institution Astrophysical Observatory		6b. OFFICE SYMBOL (If applicable) NC		7a. NAME OF MONITORING ORGANIZATION AFOSR/NC	
6c. ADDRESS (City, State, and ZIP Code) 60 Garden Street Cambridge, MA 02138		7b. ADDRESS (City, State, and ZIP Code) Building 410, Bolling AFB DC 20332-6448			
8a. NAME OF FUNDING / SPONSORING ORGANIZATION AFOSR		8b. OFFICE SYMBOL (If applicable) NC		9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER AFOSR-88-0042	
8c. ADDRESS (City, State, and ZIP Code) Building 410, Bolling AFB DC 20332-6448		10. SOURCE OF FUNDING NUMBERS			
		PROGRAM ELEMENT NO. 61102F		PROJECT NO. 2303	TASK NO. B3
				WORK UNIT 3	ACCESSION NO. 3
11. TITLE (Include Security Classification) (U) THEORETICAL INVESTIGATIONS OF METASTABILITY AND ENERGY STORAGE IN MOLECULAR SYSTEMS					
12. PERSONAL AUTHOR(S) Kate P. Kirby					
13a. TYPE OF REPORT FINAL		13b. TIME COVERED FROM 11/89 TO 11/91		14. DATE OF REPORT (Year, Month, Day) 3/9/93	
15. PAGE COUNT 12					
16. SUPPLEMENTARY NOTATION					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP	Metastability; diatomic molecules; excited states; lifetimes		
19. ABSTRACT (Continue on reverse if necessary and identify by block number)					
<p>1 A manuscript was completed describing the metastable vibrational levels of the $D^1\Delta$ and $I^1\Sigma$ states of CO. Methods of population of these levels have been investigated. Work on binding in several high-spin multiplicity systems, Σ^+ states in CN and NO, is almost complete..</p>					
20. DISTRIBUTION / AVAILABILITY OF ABSTRACT <input type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED		
22a. NAME OF RESPONSIBLE INDIVIDUAL Dr Michael R. Berman			22b. TELEPHONE (Include Area Code) (202) 767-4963		22c. OFFICE SYMBOL AFOSR/NC

THEORETICAL INVESTIGATIONS OF
METASTABILITY AND ENERGY STORAGE
IN MOLECULAR SYSTEMS

AFOSR-88-0042

Annual Technical Report

For the period 1 November 1989 through 31 October 1990

and

For the period 1 November 1990 through 31 October 1991

Principal Investigator
Dr. Kate P. Kirby

March 1993

DTIC QUALITY INSPECTED 5

Prepared for
Air Force Office of Scientific Research
Bolling AFB, D. C. 20332

Accession For	
NTIS CRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution /	
Availability Codes	
Dist	Avail and/or Special
A-1	

Smithsonian Institution
Astrophysical Observatory
Cambridge, Massachusetts 02138

The Smithsonian Astrophysical Observatory
is a member of the
Harvard-Smithsonian Center for Astrophysics

93-26591

The AFOSR Technical Officer for this grant is Major Larry Davis,
Directorate of Chemical and Atmospheric Sciences, AFOSR/NC,
Bolling AFB, D. C. 20332

95 11 ~ 060

Annual Report

The work on the metastable vibrational levels of CO has been published in Physical Review Letters 68, 3865 (1992). The manuscript is appended to this report. The discovery of the long-lived $v = 0$ levels in the $I^1\Sigma^-$ and $D^1\Delta$ states which lie $\sim 8\text{eV}$ above the ground state may have interesting consequences. One consequence might be the use of such states as stepping stones to spectroscopic regimes not probed heretofore. Another might be the possibility of observing an electric quadrupole transition from such levels.

The work on binding in high-spin states of two molecules, CN and NO, is nearly completed. A few calculations to explore possible basis set superposition error in the present results will have to be completed before the paper is written up.

15:05 6 August, 1991

The population of long-lived vibrational levels of CO: $I^1\Sigma^-$ and $D^1\Delta$

Kate Kirby⁽¹⁾, Marcy E. Rosenkrantz⁽²⁾, and David L. Cooper⁽³⁾

⁽¹⁾ Harvard-Smithsonian Center for Astrophysics,

60 Garden Street, Cambridge, MA 02138, USA

⁽²⁾ Air Force Astronautics Laboratory, Edwards Air

Force Base, CA 93523, USA

⁽³⁾ Department of Chemistry, University of Liverpool,

P. O. Box 147, Liverpool, L69 3BX, UK

The lowest vibrational energy levels of the $I^1\Sigma^-$ and $D^1\Delta$ states of CO, which lie approximately 8 eV above the $X^1\Sigma^+$ ground state, are predicted to have very long radiative lifetimes. In the case of $I^1\Sigma^-(v=0)$ the radiative lifetime for allowed electric dipole transitions is infinite and in the case of $D^1\Delta(v=0)$ the equivalent lifetime is calculated to be 1.6 s. These states cannot predissociate and the couplings with levels of the $A^1\Pi$ state are very small. There is, therefore, the possibility of using the $v=0$ levels of the $I^1\Sigma^-$ and $D^1\Delta$ states as reservoirs for highly energetic molecules and to open up spectroscopic regions which are otherwise inaccessible. It might even be possible to observe electric quadrupole or magnetic dipole transitions from these levels. In order to encourage experimental studies, methods are proposed for populating the $v=0$ levels of the $I^1\Sigma^-$ and $D^1\Delta$ states, including stimulated emission from the $W^1\Pi$ state.

The phenomenon of metastability, in which an excited state lives orders of magnitude longer than is usual before giving up its energy, is a powerful tool to be used to advantage in the laboratory. Metastable states can be used as "stepping stones" to an energy regime or to entire multiplet systems which are not easily accessible from the ground state. For example, studies of the triplet manifold in H_2 depend critically on the existence of the $c^3\Pi_u^-$ metastable state.¹⁻⁴ The absence of a single fast decay channel permits the study of competing slower processes such as magnetic dipole and electric quadrupole transitions.

Carbon monoxide is an ideal system in which to explore the phenomenon of metastability, in that it has been extensively studied spectroscopically^{5,6} and it has been isolated in inert gas matrices.⁷ In a previous paper⁸ by Rosenkrantz and Kirby (RK), calculations of the potential energy curves and electronic wavefunctions for the $I^1\Sigma^-$ and $D^1\Delta$ states of CO were reported. RK computed radiative lifetimes of different vibrational levels, considering vibrational cascade within each electronic state and also decay to the nearby $A^1\Pi$ state. In this paper we focus on the $v=0$ levels of the $I^1\Sigma^-$ and $D^1\Delta$ states, which were shown by RK to have very long radiative lifetimes. We investigate ways of populating these levels, in order to encourage experimental studies of their possible decay mechanisms.

Ab initio potential energy curves⁸⁻¹⁰ for low-lying singlet states of CO are illustrated in Figure 1. It is important to notice that the $A^1\Pi$, $I^1\Sigma^-$ and $D^1\Delta$ states all lie at approximately the same energy (~ 8 eV) above the $X^1\Sigma^+$ ground state and that there are no intervening singlet states. The $I^1\Sigma^-$ and $D^1\Delta$ states, which both arise from the electronic configuration $\dots 5\sigma^2 1\pi^3 2\pi$, have very similar potential energy curves and one-electron properties. Electric dipole transitions from the $I^1\Sigma^-$ and $D^1\Delta$ states to the ground state are forbidden, but these two excited states can, in principle, decay to the $A^1\Pi$ state. The close proximity of the $A^1\Pi$ state leads to transition probabilities from vibrational levels v' of the $I^1\Sigma^-$ and $D^1\Delta$ states which range over five orders of magnitude, decreasing significantly with decreasing v' .⁸ An analogous phenomenon has been noted¹¹ for the isoelectronic system N_2 , in which the $W^3\Delta_u$ state lies nearly coincident with the only state to which it can radiate via an allowed electric dipole transition, namely the $B^3\Pi_g$ state.

Although the minimum of the $A^1\Pi$ state of CO lies lower than that of the $I^1\Sigma^-$ state by 8.6 cm^{-1} ,¹² the difference in the shapes of these two potentials is such that the $I^1\Sigma^-(v=0)$ level actually

lies lower than $A^1\Pi(v=0)$ by $\sim 220\text{ cm}^{-1}$. The $I^1\Sigma^-(v=0)$ level cannot radiate by an allowed electric dipole transition and will thus be very long lived. The minimum of the $D^1\Delta$ state lies 852.2 cm^{-1} above that of the $A^1\Pi$ state.¹² The $D^1\Delta(v=0)$ level lies $\sim 610\text{ cm}^{-1}$ above $A^1\Pi(v=0)$, and this is the one level to which it can radiate by an allowed electric dipole transition. The corresponding lifetime is 1.6 s.

Various vibration-rotation levels of the $I^1\Sigma^-$ and $D^1\Delta$ states have been observed by Simmons, Bass and Tilford¹³ as perturbers of the upper state ($A^1\Pi$) of the fourth positive system. The $v=0$ levels have not been implicated in these perturbations. Also, due to l-uncoupling interactions with certain levels of the $A^1\Pi$ state, it has proved possible to observe very weak absorption from $X^1\Sigma^+(v''=0)$ to certain vibrational levels of the $I^1\Sigma^-$ and $D^1\Delta$ states.¹⁴ The $I^1\Sigma^-(v'=0)$ and $D^1\Delta(v'=0)$ levels have not been observed in this way. The $I^1\Sigma^-$ and $D^1\Delta$ states have also been populated in two-photon transitions from $X^1\Sigma^+(v''=0)$ by Kittrell and co-workers.¹⁵ The only levels observed in this way are $D^1\Delta(v'=7,10,12)$ and $I^1\Sigma^-(v'=7)$.

Experiments on the electron impact excitation of the $I^1\Sigma^-$ and $D^1\Delta$ states^{16,17} do not yet have vibrational resolution, but it seems clear from appearance potentials of 9.5 to 10.0 eV, and the measured lifetimes $\tau \sim 80\text{ }\mu\text{s}$ or $\tau \sim 97\text{ }\mu\text{s}$, that both sets of experiments are consistent with the population of levels with $v \geq 10$.

In spite of the range of observations of the $D^1\Delta$ and $I^1\Sigma^-$ states of CO, there is no evidence that the $v=0$ levels of either of these states has ever been observed. The fundamental problem with populating the $v'=0$ levels of the $I^1\Sigma^-$ and $D^1\Delta$ states from $X^1\Sigma^+(v''=0)$ either by two-photon absorption or by electron impact excitation is that the Franck-Condon overlaps for such a transition are vanishingly small. The Franck-Condon overlaps of the $v'=0$ and $v'=1$ levels of the $I^1\Sigma^-$ state are shown in Figure 2 as functions of v'' for the $X^1\Sigma^+$ state. The corresponding Figure for the $D^1\Delta$ state is very similar, and thus is not shown. The strongest transitions populating the $v'=0$ levels, with $|\text{overlap}|^2$ greater than 0.1, originate from ground state levels with $v''=8$ to 12. Vibrationally excited levels of the $X^1\Sigma^+$ state have been used before in absorption spectroscopy: see, for example, studies of the $B^1\Sigma^+ \leftarrow X^1\Sigma^+$ transition by Wolk and Rich¹⁸ and of $A^1\Pi \leftarrow X^1\Sigma^+$ by DeLeon.¹⁹

Another possible means of populating the $v=0$ levels of the $I^1\Sigma^-$ and $D^1\Delta$ states is by cascade from higher-lying vibrational levels of these states. However, this cascade is predicted to be much less

probable than radiative decay to the $A^1\Pi$ state.⁸ The one exception is the $v=1$ level of the $I^1\Sigma^-$ state which preferentially (by a factor of two) radiates to $I^1\Sigma^-(v=0)$. Another plausible mechanism for populating these elusive levels is charge transfer from CO^+ . Indeed, the most effective mechanism for populating the important $c^3\Pi_u^-$ state of H_2 is charge transfer collisions of H_2^+ in caesium vapour.⁴

An interesting alternative mechanism for populating $I^1\Sigma^-(v=0)$ and $D^1\Delta(v=0)$ involves emission from the higher-lying $W^1\Pi$ state. Transitions involving the $W^1\Pi$ state have been observed by Ogawa and Ogawa,²⁰ Tilford and Simmons,²¹ and by Sekine, Adachi and Hirose.²² This state is the first member of a Rydberg series converging to the $A^2\Pi$ excited state of CO^+ . Cooper and Kirby¹⁰ have carried out an *ab initio* study of this state, including a calculation of the $W^1\Pi-X^1\Sigma^+$ oscillator strength, which confirmed the very strong nature of this transition.

Using wavefunctions presented previously,⁸⁻¹⁰ we have now calculated transition probabilities for the decay of various vibrational levels of the $W^1\Pi$ state to seven lower-lying singlet electronic states. The results are summarized in Table I. Null entries in the Table denote transition probabilities which are estimated to be less than $0.05 \times 10^7 \text{ s}^{-1}$. Most of the radiative decay from the $W^1\Pi$ state takes place to the $X^1\Sigma^+$ state. The $W^1\Pi-C^1\Sigma^+$ and $W^1\Pi-E^1\Pi$ transition probabilities are particularly small because of the small v^3 factors for these transitions. Transition probabilities to the $D^1\Sigma^+$ state are negligible. Our calculations are consistent with the observation of Sekine *et al.*²² that the $W^1\Pi(v'=0)-B^1\Sigma^+(v''=0)$ transition is weak. We calculate a transition probability for this transition of $0.03 \times 10^7 \text{ s}^{-1}$.

We predict the largest $W^1\Pi(v')-X^1\Sigma^+(v''=0)$ oscillator strength to be for $v'=2$. However, no emission has ever been observed for vibrational levels of the $W^1\Pi$ state with $v' \geq 1$ and it appears that strong predissociation eradicates all but the lowest vibrational level.

The transition probabilities from the $W^1\Pi(v'=0)$ state are shown in Figure 3 as functions of v'' for the $I^1\Sigma^-$ and $D^1\Delta$ states. For $W^1\Pi(v'=0)$ we predict branching ratios for radiative decay to the $I^1\Sigma^-$ and $D^1\Delta$ states of 2% and 3%, respectively. Branching ratios for radiative decay to the $v''=0$ levels of the $I^1\Sigma^-$ and $D^1\Delta$ states are approximately 0.15% in each case. Transition probabilities to $v''=0$ and to $v'' \neq 0$ are collected in Table II. These branching ratios could be made considerably more favourable by

the process of stimulated emission from $W^1\Pi(v'=0)$, in which a laser of suitable frequency is used to populate preferentially $I^1\Sigma^-(v''=0)$ or $D^1\Delta(v''=0)$.

Finally, having suggested methods for producing these elusive vibrational levels, we speculate very briefly on possible decay mechanisms. Both the $I^1\Sigma^-$ and $D^1\Delta$ states have allowed magnetic dipole and electric quadrupole transitions to lower-lying states. Electric quadrupole transitions from $I^1\Sigma^-$ and $D^1\Delta$ to the $X^1\Sigma^+$ state appear to have lifetimes on the order of $\tau \sim 0.1$ s, assuming ΔE values on the order of 6 eV. An identification of an electric quadrupole transition for this molecule would be noteworthy, as such transitions do not appear to have been observed for heteronuclear diatomics. In addition, there is the possibility of intensity stealing from allowed electric dipole transitions. A likely scenario is based on spin-orbit interactions with the $a^3\Pi$ state, which is the upper state of the much-studied Cameron bands. The $v=0$ levels of the $I^1\Sigma^-$ and $D^1\Delta$ states are located near $v=10$ and $v=11$ of the $a^3\Pi$ state, and so any perturbations are likely to be strongly J dependent.

Of our various proposals for generating the long-lived $v'=0$ levels of the $I^1\Sigma^-$ and $D^1\Delta$ states, the most interesting one, and potentially the most useful, is stimulated emission from the $W^1\Pi$ state. It is worth recalling that this Rydberg state is connected via a very strong transition ($f_{00}=0.005$) to the $v''=0$ level of the $X^1\Sigma^+$ ground state.

Acknowledgment: This research was supported by a grant from the Air Force Office of Scientific Research, AFOSR-88-0042.

Table I.

Transition probabilities $A_{v'}$ (in 10^7 s^{-1}) for vibrational levels of the $W^1\Pi$ state decaying via allowed electric dipole transitions to lower-lying electronic states. A null entry indicates a value not greater than $0.05 \times 10^7 \text{ s}^{-1}$.

v'	$X^1\Sigma^+$	$B^1\Sigma^+$	$C^1\Sigma^+$	$A^1\Pi$	$E^1\Pi$	$I^1\Sigma^-$	$D^1\Delta$
0	23.8	-	-	0.1	-	0.5	0.8
1	23.2	0.1	-	0.2	-	0.5	0.7
2	26.7	-	-	0.1	-	0.5	0.7
3	29.3	-	-	0.1	-	0.5	0.7
4	28.9	0.1	-	0.1	-	0.5	0.7

Table II.

Transition probabilities $A_{v''v'}$ (in 10^7 s^{-1}) from given vibrational levels v' of the $W^1\Pi$ state to vibrational levels v'' of the $I^1\Sigma^-$ and $D^1\Delta$ states.

v'	$I^1\Sigma^-$		$D^1\Delta$	
	$A_{0v'}$	$\sum_{v'' \neq 0} A_{v''v'}$	$A_{0v'}$	$\sum_{v'' \neq 0} A_{v''v'}$
0	0.03	0.47	0.04	0.71
1	0.11	0.41	0.12	0.62
2	0.16	0.36	0.19	0.55
3	0.14	0.36	0.20	0.73
4	0.08	0.40	0.14	0.72

Figure captions

Figure 1

Ab initio potential energy curves for low-lying singlet states of CO.

Figure 2

Franck-Condon overlaps of the $v'=0$ (\longrightarrow) and $v'=1$ (\dashrightarrow) levels of the $I^1\Sigma^-$ state with levels v'' of the $\lambda^1\Sigma^+$ state.

Figure 3

Transition probabilities $A_{v''}$ for electric dipole transitions from $W^1\Pi(v'=0)$ to levels v'' of the $I^1\Sigma^-$ (\longrightarrow) and $D^1\Delta$ (\dashrightarrow) states.




Figure 1




Figure 2




Figure 3

References

- [1] W. Lichten, T. Wik and T.A. Miller, J. Chem. Phys. 71, 2441 (1979).
- [2] E.E. Eyler and F.M. Pipkin, Phys. Rev. Lett. 47, 1270 (1981).
- [3] H. Helm, D.P. de Bruijn and J. Los, Phys. Rev. Lett. 53, 1642 (1984).
- [4] L.J. Lembo, D.L. Huestis, S.R. Keiding, N. Bjerre and H. Helm, Phys. Rev. A 38, 3447 (1988).
- [5] S.G. Tilford and J.D. Simmons, J. Phys. Chem. Ref. Data 1, 147 (1972).
- [6] P.H. Krupenie, Nat. Stand. Ref. Data Ser., Nat. Bur. Stand. 5, 1 (1966).
- [7] J. Bahrdt and N. Schwentner, J. Chem. Phys. 88, 2869 (1988); J. Bahrdt, H. Nahme and N. Schwentner, Chem. Phys. 144, 273 (1990).
- [8] M.E. Rosenkrantz and K. Kirby, J. Chem. Phys. 90, 6528 (1989).
- [9] D.L. Cooper and K. Kirby, J. Chem. Phys. 87, 424 (1987); K. Kirby and D.L. Cooper, J. Chem. Phys. 90, 4895 (1989).
- [10] D.L. Cooper and K. Kirby, Chem. Phys. Lett. 152, 393 (1988); Erratum, *ibid.* 155, 624 (1989).
- [11] R. Covey, K.A. Saum and W. Benesch, J. Opt. Soc. Am. 63, 592 (1973).
- [12] K.P. Huber and G. Herzberg, Molecular Spectra and Molecular Structure Vol. 4, Constants of Diatomic Molecules (Van Nostrand, Princeton, 1979).
- [13] J.D. Simmons, A.M. Bass and S.G. Tilford, Astrophys. J. 155, 345 (1969).
- [14] J.D. Simmons and S.G. Tilford, J. Chem. Phys. 45, 2965 (1966); G. Herzberg, J.D. Simmons, A.M. Bass, and S.G. Tilford, Can. J. Phys. 44, 3039 (1969).
- [15] C. Kittrell, S. Cameron, L. Butler, R.W. Field, and R.F. Barrow, J. Chem. Phys. 78, 3623 (1983); B.A. Garetz and C. Kittrell, Phys. Rev. Lett. 53, 156 (1984); B.A. Garetz, C. Kittrell and A.C. LeFloch, J. Chem. Phys. 94, 843 (1991).
- [16] W.C. Wells, W.L. Borst and E.C. Zipf, Phys. Rev. A 8, 2463 (1973).
- [17] N.J. Mason and W.R. Newell, J. Phys. B: At. Mol. Opt. Phys. 21, 1293 (1988).
- [18] G. L. Wolk and J. W. Rich, J. Chem. Phys. 79, 12 (1983).
- [19] R. L. DeLeon, J. Chem. Phys. 89, 20 (1988).
- [20] M. Ogawa and S. Ogawa, J. Mol. Spectrosc. 49, 454 (1974).
- [21] S.G. Tilford and J.G. Simmons, J. Mol. Spectrosc. 53, 436 (1974).

- [22] S. Sekine, Y. Adachi and C.Hirose, J. Chem. Phys. 90, 5346 (1989).